

University of Groningen

Full ablation of C9orf72 in mice causes immune system-related pathology and neoplastic events but no motor neuron defects

Sudria-Lopez, Emma; Koppers, Max; de Wit, Marina; van der Meer, Christiaan; Westeneng, Henk-Jan; Zundel, Caroline A. C.; Youssef, Sameh A.; Harkema, Liesbeth; de Bruin, Alain; Veldink, Jan H.

Published in:
Acta Neuropathologica

DOI:
[10.1007/s00401-016-1581-x](https://doi.org/10.1007/s00401-016-1581-x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Sudria-Lopez, E., Koppers, M., de Wit, M., van der Meer, C., Westeneng, H.-J., Zundel, C. A. C., Youssef, S. A., Harkema, L., de Bruin, A., Veldink, J. H., van den Berg, L. H., & Pasterkamp, R. J. (2016). Full ablation of C9orf72 in mice causes immune system-related pathology and neoplastic events but no motor neuron defects. *Acta Neuropathologica*, 132(1), 145-147. <https://doi.org/10.1007/s00401-016-1581-x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CORRESPONDENCE

Full ablation of *C9orf72* in mice causes immune system-related pathology and neoplastic events but no motor neuron defects

Emma Sudria-Lopez¹ · Max Koppers^{1,2} · Marina de Wit¹ · Christiaan van der Meer¹ ·
Henk-Jan Westeneng² · Caroline A. C. Zundel¹ · Sameh A. Youssef³ ·
Liesbeth Harkema^{3,4} · Alain de Bruin^{3,4} · Jan H. Veldink² · Leonard H. van den Berg² ·
R. Jeroen Pasterkamp¹

Received: 22 April 2016 / Revised: 11 May 2016 / Accepted: 11 May 2016 / Published online: 20 May 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Non-coding hexanucleotide (GGGGCC) repeat expansions in *C9ORF72* are the most common genetic cause of amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD; C9ALS/FTD). Decreased *C9orf72* protein levels in C9ALS/FTD patients [4] support the idea that *C9ORF72* haploinsufficiency may contribute to disease pathogenesis. To test this hypothesis, we previously generated and analyzed neural-specific *C9orf72* knockout mice. Our results showed that neural-specific ablation of *C9orf72* (3110043O21Rik) in mice does not cause motor neuron degeneration or changes in motor function [3]. We therefore concluded that loss of *C9ORF72* on its own is unlikely to cause ALS and that reducing *C9ORF72* levels may comprise a promising strategy to treat C9-ALS patients. This therapeutic potential led us, and others [1, 2], to subsequently analyze knockout mice lacking *C9orf72* in all tissues. Importantly, in contrast to our previous report, we find that full ablation of *C9orf72* induces reduced survival

(Fig. 1a), which is in line with a recent study by Atanasio et al. [1] who report, but do not specify, decreased survival rates. In line with our previous observations [3], full *C9orf72* ablation results in a 5.9 % decrease in body weight ($P = 0.0056$), without affecting motor function (accelerating rotarod and grip strength test) or inducing pathological hallmarks of ALS (see also [1, 2]), such as motor neuron degeneration, gliosis, enhanced ubiquitination and TDP-43 mislocalization. However, post-mortem analysis of full *C9orf72* knockout mice ($n = 5$; 11–15 months of age) revealed enlarged lymph nodes (LNs) ($n = 4$ mice) and splenomegaly ($n = 5$) (Fig. 1b). Detailed histological evaluation detected massive infiltration of histiocytes/macrophages and lymphocytes in multiple organs, including LNs, spleen, bone marrow, liver, kidney and lung (Fig. 1c–k). In addition to these immunological phenotypes, which are in part also reported by Atanasio et al. [1] and O'Rourke et al. [2], we detect evidence of neoplastic events. LNs of several animals ($n = 4$) contained infiltrates of B220/CD45R-positive B-lymphocytes that disrupted tissue architecture and were accompanied by increased expression of the proliferation marker Ki67, suggesting the development of B-cell lymphomas (Fig. 1c–e). Furthermore, disrupted tissue architecture and homogeneous populations of F4/80-positive macrophages expressing Ki67 were present in LNs ($n = 3$), spleen ($n = 3$), liver ($n = 1$) and lung ($n = 1$). Moreover, infiltrating cells in the liver and lung accumulated in intravascular spaces (Fig. 1f–k), suggesting the occurrence of metastatic histiocytic sarcomas. These results indicate that the defects in immune cell function recently reported in *C9orf72* knockout mice (e.g. changes in endosome/lysosomal trafficking, cytokine production) [1, 2] may ultimately lead to neoplastic events in multiple organs. These findings have important implications as they indicate that strategies aimed at lowering systemic *C9ORF72* levels

L. H. van den Berg and R. J. Pasterkamp are joint senior authors.

✉ R. Jeroen Pasterkamp
r.j.pasterkamp@umcutrecht.nl

- ¹ Department of Translational Neuroscience, Brain Center Rudolf Magnus, University Medical Center Utrecht, Universiteitsweg 100, 3584 CG Utrecht, The Netherlands
- ² Department of Neurology and Neurosurgery, Brain Center Rudolf Magnus, University Medical Center Utrecht, 3584 CX Utrecht, The Netherlands
- ³ Department of Pathobiology, Faculty of Veterinary Medicine, Dutch Molecular Pathology Center, Utrecht University, 3584 CL Utrecht, The Netherlands
- ⁴ Division of Molecular Genetics, Department of Pediatrics, University Medical Center Groningen, 9713 AV Groningen, The Netherlands




Fig. 1 *C9orf72* knockout mice display reduced survival, immune system-related pathology and neoplastic events. **a** Kaplan–Meier curves show survival rates corrected for gender and body weight. *C9orf72* knockout mice show reduced survival as compared to littermate controls (Hazard ratio = 19.0; 95 %, CI: 2.4–150.2, $P = 0.005$). Wild-type controls $n = 24$; *C9orf72* knockout $n = 29$. **b** Gross image showing enlarged lymph nodes (LNs; *black arrows*) and splenomegaly (*white arrow*) in a *C9orf72* knockout mouse (12 months of age). **c–e** B-cell lymphoma in the submandibular LNs of *C9orf72* knockout mouse. Nodal tissue is effaced by a monotypic cell population composed of B220/CD45R-positive lymphocytes (B cells). Note

the high proliferation rate of the neoplastic lymphocytes as indicated by immunostaining for Ki67 (proliferation marker). **f–h** Histiocytic sarcoma in the liver of *C9orf72* knockout mouse. Hepatic sinusoids are filled with atypical histiocytes and multinucleated giant cells that stain positive for the macrophage lineage marker F4/80 and exhibit a high proliferation rate, as evidenced by Ki67 immunostaining. **i–k** Histiocytic sarcoma in lung vasculature in *C9orf72* knockout mouse. Pulmonary blood vessels are filled with atypical and multinucleated giant cells that immunostain for F4/80 and Ki67. *H&E* hematoxylin and eosin. Scale bar 1.3 cm (**b**), 65 μ m (**c**), 40 μ m (**d–h**), 125 μ m (**i**), and 90 μ m (**j, k**)

in C9ALS/FTD patients may have negative side effects and that emphasis should be on therapeutic approaches that selectively target the hexanucleotide repeat expansions or their downstream pathogenic effects.

Acknowledgments This study was supported by funding from the Netherlands Organization for Health Research and Development (NWO-VICI; LvdB and RJP); Thierry Latran Foundation (JHV and RJP), Prinses Beatrix Spierfonds (LvdB and RJP), Van Meer Stichting, and Netherlands ALS foundation (TOTALS; LvdB and RJP).

Compliance with ethical standards

Conflict of interest The authors declare no conflicts of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Atanasio A, Decman V, White D et al (2016) C9orf72 ablation causes immune dysregulation characterized by leukocyte expansion, autoantibody production, and glomerulonephropathy in mice. *Sci Rep* 6:23204. doi:[10.1038/srep23204](https://doi.org/10.1038/srep23204)
2. O'Rourke JG, Bogdanik L, Yáñez A et al (2016) C9orf72 is required for proper macrophage and microglial function in mice. *Science* 351:1324–1329. doi:[10.1126/science.aaf1064](https://doi.org/10.1126/science.aaf1064)
3. Koppers M, Blokhuis AM, Westeneng HJ et al (2015) C9orf72 ablation in mice does not cause motor neuron degeneration or motor deficits. *Ann Neurol* 78:426–438. doi:[10.1002/ana.24453](https://doi.org/10.1002/ana.24453)
4. Waite AJ, Baumer D, East S et al (2014) Reduced C9orf72 protein levels in frontal cortex of amyotrophic lateral sclerosis and frontotemporal degeneration brain with the C9ORF72 hexanucleotide repeat expansion. *Neurobiol Aging* 35:1779.e5–1779.e13. doi:[10.1016/j.neurobiolaging.2014.01.016](https://doi.org/10.1016/j.neurobiolaging.2014.01.016)